

Prior Emission Model for X-ray Plateau Phase of Gamma-Ray Burst Afterglows

Ryo Yamazaki¹

Department of Physical Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan

ryo@theo.phys.sci.hiroshima-u.ac.jp

ABSTRACT

The two-component emission model to explain the plateau phase of the X-ray afterglows of gamma-ray bursts (GRBs) is proposed. One component, which is responsible for the plateau and subsequent normal decay phase of the X-ray afterglow, is the prior emission via outflow ejected from the central engine before the main burst. The other is the main outflow, which causes the prompt GRB emission and the initial steep decay phase of the X-ray afterglow. In this model, the transition from the plateau to the subsequent normal decay phase is an artifact of the choice of the zero of time. For events with distinct plateau phase, the central engine is active 10^3 – 10^4 sec before the launch of the main outflow. According to this model, a prior emission in the X-ray and/or optical bands 10^3 – 10^4 sec before the prompt GRB emission is possibly seen, which will be tested by near-future instruments such as Monitor of All-sky X-ray Image (MAXI), WIDE-field telescope for GRB Early Timing (WIDGET), and so on.

Subject headings: gamma rays: bursts —gamma rays: theory

1. Introduction

The X-Ray Telescope (XRT) onboard *Swift* has revealed complex temporal behavior of the X-ray afterglows of gamma-ray bursts (GRBs) in the first few hours (Burrows et al. 2005; Tagliaferri et al. 2005; Nousek et al. 2006; O’Brien et al. 2006; Willingale et al. 2007; Zhang et al. 2007; Liang et al. 2007, 2008). This time window had been largely unexplored before the *Swift* era, and studies of early afterglows have revealed many questions concerning

GRBs, such as the emission mechanism, nature of the central engine, and burst environment (e.g., Zhang et al. 2006; Zhang 2007).

Early X-ray afterglows have three phases¹, which was not predicted by the standard model from the pre-*Swift* era (Nousek et al. 2006; Zhang 2007). *Phase I: initial steep decay phase*. Initially, the X-ray afterglow decays very steeply; the most popular interpretation is that this is the tail emission of the prompt GRB (Kumar & Panaitescu 2000; Zhang et al. 2006; Yamazaki et al. 2006; Zhang et al. 2009), although other possibilities have been proposed (e.g., Zhang et al. 2007). *Phase II: plateau phase*. At several hundreds of seconds after the burst trigger, this phase begins until $\sim 10^3$ – 10^4 sec, whose origin is quite uncertain. This is the main topic of this Letter. *Phase III: normal decay phase*. After the plateau phase ends, the X-rays subsequently decay with the decay index usually steeper than unity, as expected in the pre-*Swift* era. This decay behavior is well explained by the classical external shock model (Sari et al. 1998), in which neither the delayed energy injection nor the time dependency of shock microphysics is considered.

Phase II is the most enigmatic in early X-ray afterglows. So far, various kinds of models have been proposed such as the energy injection model (Nousek et al. 2006; Zhang et al. 2006; Granot & Kumar 2006), the inhomogeneous or two-component jet model (Toma et al. 2006; Eichler & Granot 2006; Granot et al. 2006), the time-dependent microphysics model (Ioka et al. 2006; Granot et al. 2006; Fan & Piran 2006), the reverse shock-dominated afterglow model (Genet et al. 2007), the prior activity model (Ioka et al. 2006), the internal engine model (Ghisellini et al. 2007), the cannonball model (Dado et al. 2006), and the dust scattering model (Shao & Dai 2007). In this Letter, another model of phase II is proposed.

2. Two-component Emission Model

As described in the following, we consider a two-component emission model in which a prior and the main outflows emit X-rays independently (see fig. 1). One component is the prior emission via outflow ejected from the central engine before the main burst. This is responsible for phases II and III of the X-ray afterglow. This emission component arrives at the observer before the main burst triggering prompt GRB detectors (i.e., the onset of GRB) such as BAT onboard *Swift*. It decays with time simply in a single power-law form

$$f_0(t) = A_0 t^{-\alpha_0} \quad , \quad (1)$$

¹ For simplicity, the X-ray flares are not considered in this Letter.

where A_0 is a constant and the time coordinate, t , is measured in the rest frame of the observer. The epoch $t = 0$ is taken around the time of arrival at the observer of the (unseen) information of the launch of the outflow at the central engine. This kind of a choice of the time zero is seen in many references (e.g., Panaitescu & Kumar 2000; Kobayashi & Zhang 2007). The origin of the prior emission, f_0 , is not discussed in detail here. It can be either internal engine activity or external shock emission of the outflow.

We set an observer time, T , where $T = 0$ corresponds to the onset of the prompt GRB. The interval between the time $t = 0$ and $T = 0$ is assumed to be T_0 seconds, that is, $t = T + T_0$. Then, one obtains

$$f_0(T) = A_0(T + T_0)^{-\alpha_0} \quad , \quad (2)$$

which becomes constant if $T \ll T_0$, while $f_0 \propto T^{-\alpha_0}$ when $T \gg T_0$. In order to explain phases II and III of the X-ray afterglow, T_0 should be 10^3 – 10^4 sec, and α_0 should be the temporal decay index of phase III. It is noted that the onset time of the prior emission is unknown. The first detectable X-rays from the prior component arrive at the observer in the time range $0 \lesssim t \lesssim T_0$ ($-T_0 \lesssim T \lesssim 0$). This fact will be further discussed in § 3.5. Another remark is that from Eq. (2) alone, the introduction of T_0 shifts the origin of time. It has been known that, for phase I, the choice of time zero affects the decay slope (Zhang et al. 2006; Yamazaki et al. 2006). The same argument for phase II is used for the first time in this Letter.

The other component is the main outflow, which causes the prompt GRB emission and the subsequent phase I of the X-ray afterglow. With time coordinate T , phase I of the X-ray afterglow is well approximated by a single power-law model given by

$$f_1(T) = A_1 T^{-\alpha_1} \quad , \quad (3)$$

where A_1 is a constant and $\alpha_1 \approx 3 - 6$.

The whole light curve of the X-ray afterglow from phase I to III is described by the sum of the two components introduced above, that is,

$$\begin{aligned} f(T) &= f_0(T) + f_1(T) \\ &= A_0(T + T_0)^{-\alpha_0} + A_1 T^{-\alpha_1} \quad . \end{aligned} \quad (4)$$

This is our formula for observed X-ray afterglows, where $f_0(T)$ describes phases II and III, while $f_1(T)$ fits phase I.

We find that by choosing appropriate values of the parameters, α_0 , α_1 , T_0 , A_0 , and A_1 , the observed light curves of X-ray afterglows are well described with Eq. (4). Figure 2 shows some examples of the fit.

It is not surprising that our formula, Eq. (4), well explains the observational results of X-ray afterglows. The functional form of Eq. (2) is a good approximation of that introduced by Willingale et al. (2007):

$$f_a(T) = \begin{cases} F_a \exp[\alpha_a(1 - T/T_a)] \exp(-t_a/T) , & T < T_a , \\ F_a (T/T_a)^{-\alpha_a} \exp(-t_a/T) , & T_a \leq T . \end{cases} \quad (5)$$

They have shown that the observed light curves of phases II and III are well fitted with Eq. (5). The function $f_a(T)$ becomes constant with T if $t_a \ll T \ll T_a$, while $f_a \propto T^{-\alpha_a}$ if $T_a \ll T$. This behavior is quite similar to that of $f_0(T)$. Indeed, one can find that over a wide parameter range, $f_a(T) \approx f_0(T)$ for $t_a \ll T$ if we take $t_a \ll T_a \approx T_0$ and $\alpha_a \approx \alpha_0$.

3. Discussion

3.1. Overall shape of the X-ray afterglow

In this Letter, we have seen that for most events, phases II and III of the observed light curves of the X-ray afterglow can be well fitted with a very simple formula, Eq. (2). In particular, the observational facts are that for most events, the transition from phase II to III is slow and fairly smooth, and that the X-ray spectrum remains unchanged across the transition (Nousek et al. 2006); this can be naturally explained by our model because the transition from phase II to III is an artifact of the choice of the time zero. A typical exception is GRB 070110; at the end of the plateau phase II, the light curve shows an abrupt drop (Troja et al. 2007). Such events may need other explanations (e.g., Kumar et al. 2008).

After phase III, subsequent fourth phase is occasionally observed: the so called *post jet break phase* (Phase IV; Zhang 2007). Its typical decay index is ~ -2 , satisfying the predictions of the jetted afterglow model (Sari et al. 1999). This phase IV can also be clearly seen in our sample (e.g., GRB 060428A; upper right of Fig. 2). If the prior emission is from an external shock of the prior outflow, the jetted afterglow model is applied to the present case. As will be discussed in § 3.4, the origin of the optical afterglow is different from the X-ray one, which explains the fact that the epoch of the jet break in the X-ray band is not generally the same as the optical one (Sato et al. 2007; Liang et al. 2008).

3.2. Observed $f_0(T_0) - T_0$ Correlation

Sato et al. (2008) investigated the characteristics of the transition from phase II to III for 11 events with known redshifts. They derived the transition time, T_{brk}^0 , and the

isotropic luminosity at that time $L_{X,\text{end}}$, and found that $L_{X,\text{end}}$ is well correlated with T_{brk}^0 as $L_{X,\text{end}} \propto (T_{\text{brk}}^0)^{-1.4}$. They adopt a broken power-law form to fit the light curve in phases II and III which is different from that considered in this Letter. However, we expect that their T_{brk}^0 and $L_{X,\text{end}}$ roughly correspond to T_0 and $f_0(T_0)$, respectively, that are considered in § 2. Indeed, one finds $f_0(T_0) \propto T_0^{-\alpha_0}$ from Eq. (2). Hence, if $\alpha_0 \approx 1.4$ which is a typical number for the decay index of phase III, we can reproduce the observed result of Sato et al. (2008).

A similar analysis has been done but with Eq. (5) as the fitting formula (Dainotti et al. 2008). For 32 events with measured redshifts, they found a correlation between T_a and the X-ray luminosity at the time T_a , $L_X(T_a)$, as $L_X(T_a) \propto T_a^{-\beta}$ with $\beta = 0.6\text{--}0.74$. Since $T_a \approx T_0$ and $L_X(T_a) \approx f_0(T_0)$, their correlation indicates $f_0(T_0) \propto T_0^{-\beta}$. The index β is smaller than the typical value of α_0 ($\approx 1.0\text{--}1.5$). However, the claimed correlation has large scatter (see Figures 1 and 2 of Dainotti et al. 2008), which may be explained by the scatter of α_0 in our model.

There is a difference between the results of Sato et al. (2008) and Dainotti et al. (2008). At present, the number of events with known redshifts may be small, so this discrepancy may be resolved if the number of events increases.

3.3. Link between T_0 and the Prompt Emission Properties

Nava et al. (2007) studied the properties of prompt emission and X-ray afterglows of 23 GRBs with known redshifts. They adopted Eq. (5) in fitting phases II and III of the X-ray afterglow, and found that for events with measured spectral peak energy E_p , the time T_a weakly correlates with the isotropic equivalent energy $E_{\gamma,\text{iso}}$ of the prompt GRB emission. One can find from Fig. 6 of Nava et al. (2007), that T_a seems to be roughly proportional to $E_{\gamma,\text{iso}}$. At present, this correlation is not firmly established because as noted by Nava et al. (2007), there are no correlation between T_a and the isotropic equivalent energy of prompt GRBs in the 15-150 keV band for a larger sample of GRBs with known redshift but unknown E_p (hence without k -correction).

The quantity $E_{\gamma,\text{iso}}$ correlates with E_p (Amati et al. 2002). Hence, if the T_a – $E_{\gamma,\text{iso}}$ correlation exists, the bright GRBs with large $E_{\gamma,\text{iso}}$ and E_p have large T_a , which is responsible for the distinct plateau phase. On the other hand, the X-ray flashes or X-ray-rich GRBs (e.g., Heise et al. 2001; Barraud et al. 2003; Sakamoto et al. 2008) have small T_a ($\approx T_0$), and have X-ray afterglow without phase II. This tendency could have been seen in Sakamoto et al. (2008).

The above arguments may lead a link between long GRBs and X-ray flashes/X-ray-rich

GRBs. Suppose that the outflow ejection is not continuous but intermittent, i.e., the central engine ejects two distinct outflows with a time interval of $\sim T_0$. Just after the launch of the prior outflow, the central engine does not have enough energy for another outflow, so that it needs to store an additional one. During the time interval $\sim T_0$, matter surrounding the central engine is accreted, increasing the gravitational binding energy. This energy is released as the main outflow causing the prompt GRB. It is expected that the larger is T_0 , the larger is the stored gravitational energy, resulting in a brighter burst with large $E_{\gamma, \text{iso}}$. On the other hand, if T_0 is small the energy of the main outflow becomes small; this is responsible for the X-ray flash or X-ray-rich GRBs. Further details will be discussed in the near future.

3.4. External Shock Emission from the Main Outflow

The main outflow that is responsible for the prompt GRB might cause external shock X-ray emission, $f_{\text{X,ext}}(T)$. In the present two-component emission model, however, $f_{\text{X,ext}}(T)$ must be dimmer than the prior X-ray emission $f_0(T)$ throughout phases II and III. Let us consider the simplest model of external shock emission of the main outflow. The relativistically expanding shell with energy E_K interacts with the surrounding medium with uniform density n_0^2 , and emits synchrotron radiation with microphysics parameters at the shock, p , ε_e , and ε_B (Sari et al. 1998). Then, in the case of slow cooling and $\nu_e < \nu_X$, the X-ray light curves are analytically calculated as $f_{\text{X,ext}}(T) \propto E_K^{(p+2)/4} \varepsilon_e^{p-1} \varepsilon_B^{(p-2)/4} T^{(2-3p)/4} \nu_X^{-p/2}$, which is independent of n_0 (e.g., Panaitescu & Kumar 2000). If $p \approx 2$, then $f_{\text{X,ext}}(T)$ hardly depends on ε_B . Before the *Swift* era, typical values had been $E_K \sim 10^{52}\text{--}10^{53}$ erg, $\varepsilon_e \sim 10^{-1}$, and $\varepsilon_B \sim 10^{-2}$ so that the external shock emission reproduced the observed late-time X-ray afterglow. In the present case, $f_{\text{X,ext}}(T)$ must be dim, which implies small E_K and/or ε_e . A similar discussion was made by Ghisellini et al. (2007, 2008). In some models, such as the energy injection model and the inhomogeneous jet model, prompt GRB emission needs high radiation efficiency, which is defined by $\varepsilon_\gamma = E_{\gamma, \text{iso}} / (E_{\gamma, \text{iso}} + E_K)$, because E_K is small at the epoch of the prompt GRB emission (Fan & Piran 2006; Granot et al. 2006; Ioka et al. 2006; Zhang et al. 2007). As discussed here, $f_{\text{X,ext}}(T)$ can be dim if ε_e is small while E_K remains large, $\sim 10^{52}\text{--}10^{53}$ erg. Hence, the present model could avoid a serious efficiency problem. On the other hand, if E_K of the outflow is small, the efficiency ε_γ should be high. Then the mechanism of prompt GRB emission is unlike a classical internal shock model (e.g.,

² Since the prior outflow may modify the circumburst medium density profile, the external shock emission from the main outflow deviates from the case of a uniform density profile. Nevertheless, we adopt the uniform density model here for the simplicity.

Thompson et al. 2007; Ioka et al. 2007).

The observed optical afterglow comes mainly from the prior outflow. For some events, the rising part of the early optical afterglow proceeds until $T \sim 10^2$ sec (Molinari et al. 2007), which is difficult to explain with prior emission. In the case of prior emission, the time zero would be shifted $T_0 \sim 10^3$ – 10^4 sec before the burst trigger, making the light curve extraordinarily spiky. Furthermore, in most cases, the transition from phase II to III is chromatic, i.e., the optical light curves do not show any break at that epoch (Panaitescu et al. 2006), although there exist a few exceptions (Liang et al. 2007; Grupe et al. 2007; Mangano et al. 2007). Hence, at least in the early epoch, the observed optical afterglow arises from the main outflow component or others, most likely an external shock emission. Indeed, there have been some observational facts that indicate different origins of X-ray and optical afterglows (e.g., Oates et al. 2007; Sato et al. 2007; Urata et al. 2007)³.

3.5. Predicted Precursor Emission?

A possible prediction of the present model is a bright X-ray precursor before the prompt GRB emission. Let us assume that the prior X-ray emission starts at $t \sim 10^2$ sec, although its onset time is fairly uncertain (see § 2). Then, from Eq. (1), the X-ray flux in the 2–10 keV band is estimated as $f_0(t) \sim 4 \times 10^{-9}(t/10^2 \text{ s})^{-1.2} \text{ erg s}^{-1} \text{ cm}^{-2}$, where $\alpha_0 \approx 1.2$ is taken as a typical value and the flux normalization constant A_0 is determined so that $f_0 \sim 1 \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ at $t \sim 10^5$ sec (Gendre et al 2008). Such emission will be detected by Monitor of All-sky X-ray Image (MAXI)⁴. The expected event rate should be a few events per year (Suzuki et al. 2008).

However, if the emission starts with $t \sim 10^2$ s, the predicted flux might be large enough to be detected by current instruments like BAT onboard *Swift*. In order to avoid this problem, the starting time of the emission should be comparable to or later than $t \sim 10^3$ sec so that the peak flux is smaller than the detection limit of the prompt GRB emission monitors. One possibility is off-axis jet emission in the context of the external shock model (Granot & Kumar 2003; Granot 2005). Due to the relativistic beaming effect, the observed X-ray emission is dim as long as the bulk Lorentz factor of the emitting outflow is larger than the inverse of the angle between the emitting matter and the observer’s line of sight. This

³ One may expect that the optical emission also arises from the prior outflow. However, it may be outshone by the main outflow component at least in the early epoch. This condition will constrain the mechanisms of the prior X-ray emission as well as the optical one.

⁴ <http://www-maxi.tksk.jaxa.jp/indexe.html>

effect shifts the peak time of the X-ray emission toward the later epoch. For an appropriate choice of parameters, we may adjust the starting time (the peak time) of the X-ray emission.

A signal of the onset of the prior emission might also be seen in the optical band. If the prior emission is extremal shock origin, the reverse shock emission might cause a bright optical flash (Sari & Piran 1999). So far, for some events, WIDGET⁵ has given observational upper limit on the prior optical emission, $V > 10$ mag, about 750 seconds before the prompt emission (e.g., Abe et al. 2006). Further observations will constrain the model parameters. In summary, in order to test the model presented in this Letter, the search for a signal in the data of sky monitors in the optical and X-ray bands is crucial.

R.Y. would like to thank Takashi Nakamura, Kunihito Ioka, Takanori Sakamoto, Atsumasa Yoshida, Motoko Suzuki, Takeshi Uehara, and the anonymous referee for useful comments and discussions. This work was supported in part by grant-in-aid from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, No. 18740153, No. 19047004.

REFERENCES

- Abe, K. et al. 2006, GCN 4931
- Amati, L. et al. 2002, A&A, 390, 81
- Barraud, C. et al. 2003, A&A, 400, 1021
- Burrows, D. N. et al. 2005, Science, 309, 1833
- Dado, S., Dar, A., & De Rujula, A. 2006, ApJ, 646, L21
- Dainotti, M. G., Cardone, V. F., & Capozziello, S. 2008, MNRAS, 391, L79
- Eichler, D. & Granot, J. 2006, ApJ, 641, L5
- Evans, P. A. et al. 2007, A&A, 469, 379
- Fan, Y. Z. & Piran, T. 2006, MNRAS, 369, 197
- Gendre, B. et al. 2008, ApJ, 683, 620

⁵ <http://cosmic.riken.go.jp/grb/widget/>

- Genet, F. et al. 2007, MNRAS, 381, 732
- Ghisellini, G. et al. 2007, ApJ, 658, L75
- Ghisellini, G. et al. 2008, MNRAS, in press (arXiv:0811.1038)
- Granot, J. 2005, ApJ, 631, 1022
- Granot, J. & Kumar, P. 2003, ApJ, 591, 1086
- Granot, J. & Kumar, P. 2006, MNRAS, 366, L13
- Granot, J. et al. 2006, MNRAS, 370, 1946
- Grupe, D. et al. 2007, ApJ, 662, 443
- Heise, J., in 't Zand, J., Kippen, R. M., & Woods, P. M. 2001, in Proc. Second Rome Workshop: Gamma-Ray Bursts in the Afterglow Era, ed. E. Costa, F. Frontera, & J. Hjorth (Berlin: Springer), 16
- Ioka, K. et al. 2006, A&A, 458, 7
- Ioka, K. et al. 2007, ApJ, 670, L77
- Kobayashi, S. & Zhang, B. 2007, ApJ, 655, 973
- Kumar, P., Narayan, R., & Johnson, J. L. 2008, Science, 321, 376
- Kumar, P., & Panaitescu, A. 2000, ApJ, 541, L51
- Liang, E. W., Zhang, B. B., Zhang, B. 2007, ApJ, 670, 565
- Liang, E. W. et al. 2008, ApJ, 675, 528
- Mangano, V. et al. 2007, A&A, 470, 105
- Molinari, E. et al. 2007, A&A, 469, L13
- Nava, L. et al. 2007, MNRAS, 377, 1464
- Nousek, J. A. et al. 2006, ApJ, 642, 389
- Oates, S. R. et al. 2007, MNRAS, 380, 270
- O'Brien, P. T. et al. 2006, ApJ, 647, 1213
- Panaitescu, A. & Kumar, P. 2000, ApJ, 543, 66

- Panaitescu, A. et al. 2006, MNRAS, 369, 2059
- Sakamoto, T. et al. 2008, ApJ, 679, 570
- Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
- Sari, R., Piran, T., & Halpern, P. 1999, ApJ, 519, L17
- Sari, R. & Piran, T. 1999, ApJ, 520, 641
- Sato, G. et al. 2007, ApJ, 657, 359
- Sato, R. et al. 2008, arXiv:0711.0903
- Shao, L. & Dai, Z. G. 2007, ApJ, 660, 1319
- Suzuki, M. et al. 2008, to appear in 3rd international MAXI workshop proceedings
- Tagliaferri, G. et al. 2005, Nature, 436, 985
- Thompson, C. et al. 2007, ApJ, 666, 1012
- Toma, K. et al. 2006, ApJ, 640, L139
- Troja, E. et al. 2007, ApJ, 665, 599
- Urata, Y., et al. 2007, ApJ, 668, L95
- Willingale, R. et al. 2007, ApJ, 662, 1093
- Yamazaki, R. et al. 2006, MNRAS, 369, 311
- Zhang, B. et al. 2006, ApJ, 642, 354
- Zhang, B. et al. 2007, ApJ, 655, 989
- Zhang, B. 2007, Chin. J. Astron. Astrophys. 7, 1
- Zhang, B. B., E. W. Liang, & Zhang, B. 2007, ApJ, 666, 1002
- Zhang, B. B. et al. 2009, ApJ, 690, L10




Fig. 1.— Schematic diagram of the model presented in this Letter. In terms of the coordinate t , the prior emission component $f_0(t)$ takes a single power-law form throughout the burst, but in terms of $T(= t - T_0)$, the function $f_0(T)$ has an artificial plateau phase (see the text for details).




Fig. 2.— Comparison of the observed light curves of X-ray afterglows with Eq. (4), where all phases I, II, and III are well described. The adopted parameters are $(\alpha_0, \alpha_1, T_0) = (1.2, 7.0, 8000 \text{ sec}), (1.0, 4.0, 5000 \text{ sec}), (1.4, 4.0, 7000 \text{ sec}),$ and $(1.6, 5.5, 4000 \text{ sec})$, for GRB 051016B, 060428A, 060814, and 061121, respectively. Data of X-ray afterglows are taken from the *Swift* online repository (Evans et al. 2007).